

Drug : **ROBAXIN TAB**

**What is this medicine?**

METHOCARBAMOL (meth oh KAR ba mole) helps to relieve pain and stiffness in muscles caused by strains, sprains, or other injury to your muscles.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: -kidney disease -seizures -an unusual or allergic reaction to methocarbamol, other medicines, foods, dyes, or preservatives -pregnant or trying to get pregnant -breast-feeding

**How should I use this medicine?**

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. Take your medicine at regular intervals. Do not take your medicine more often than directed. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.**

**NOTE: This medicine is only for you. Do not share this medicine with others.**

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only the next dose. Do not take double or extra doses.

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 20 and 25 degrees C (68 and 77 degrees F). Keep container tightly closed. Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I watch for while using this medicine?**

Tell your doctor or health care professional if your symptoms do not start to get better or if they get worse. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks. If you are taking another medicine that also causes drowsiness, you may have more side effects. Give your health care provider a list of all medicines you use. Your doctor will tell you how much medicine to take. Do not take more medicine than directed. Call emergency for help if you have problems breathing or unusual sleepiness.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: -allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue -breathing problems -confusion -seizures -unusually weak or tired Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): -dizziness -headache -metallic taste -tiredness -upset stomach

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088